

(INSERT HEALTH CARE AGENCY)

MEDICAL DIRECTIVE: Spikevax™ Bivalent (Moderna) Vaccine

Original Date: September 2022

Next Review Date: September 2023

Contact Officer: ((INSERT AGENCY CONTACT OFFICER)

Order/Delegated Procedure

Administer injectable Spikevax COVID-19 immunizations (Moderna).

Spikevax™ Bivalent (Moderna):

Individuals ≥ 18 Years of Age

- Spikevax™ Bivalent (Moderna) is authorized for use as a booster dose of 50 mcg each for those 18 years of age and older.¹
- NACI recommends that the authorized dose of a bivalent Omicron-containing mRNA COVID-19 vaccine should be offered as a booster dose to the authorized age groups (≥18 years of age). If the bivalent Omicron-containing mRNA COVID-19 vaccine is not readily available, an original mRNA COVID-19 vaccine should be offered to ensure timely protection.¹¹

Authorized Implementers

Health care providers of (ABOVE NAMED OF AGENCY) are described as: (AGENCY HEALTH CARE PROVIDERS) and who have completed the required orientation, including a review of related policies, educations training at clinics and observation by the (AGENCY DESIGNATE).

Vaccine Administration Errors and Deviations

Any deviations in this medical directive shall be documented by the Authorized Implementer and the Clinic Lead Nurse after informing the Contact officer or designate by phone. The Contact officer will notify the Authorizing Physician. *

*Note: Common errors and deviations in COVID-19 vaccine administration that are described in the Government of Canada [COVID-19 vaccine guide for youth and adults \(12 years and over\): Managing COVID-19 vaccine administration errors or deviations](#) (Last date modified: 2022-08-11 or as current)⁸ do not require notification of the Authorizing Physician, except in all cases where a vaccine is administered to a recipient in whom the vaccine is contraindicated or not indicated. For example, vaccine administered to an individual under the age for which the vaccine is authorized in Canada must be reported to the Authorizing Physician, whereas

vaccination with a partial dose of vaccine in error to a person for whom the full vaccine dose is indicated that is addressed according to the Provincial guidance does NOT require notification of the Authorizing Physician.

Indications

The Spikevax™ Bivalent (Moderna) is indicated as a booster dose for active immunization against coronavirus disease 2019 (COVID-19) caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus in individuals 18 years of age and older.¹

Contraindications

Clients are **NOT ELIGIBLE** for Spikevax™ Bivalent (Moderna) under this medical directive if there is a risk of serious adverse reaction identified during their health assessment:

- A history of anaphylactic reaction to a previous dose of the Spikevax™ Bivalent (Moderna).¹
- Individuals who are hypersensitive to the active ingredient or to any ingredients in the formulation, including any non-medicinal ingredient, or component of the container.¹ Components of the Spikevax™ Bivalent (Moderna) are listed in Appendix A

Individuals with known allergies to components of the vaccines may speak with an appropriate physician or nurse practitioner (NP) for evaluation. This assessment will enable the development of a vaccination care plan which may include receiving the vaccine under the supervision of your physician.¹⁰

Documentation of the discussion with the physician/NP may be provided to the clinic and can include a vaccination care plan and parameters the clinic should meet to provide safe vaccine administration.¹⁰

Precautions

- Individuals should seek medical attention if they develop symptoms of Bell's palsy following receipt of mRNA COVID-19 vaccines. Very rare cases of Bell's palsy (typically temporary weakness or paralysis on one side of the face) have been reported following vaccination with COVID-19 mRNA vaccines (Pfizer-BioNTech or Moderna) in Canada and internationally among individuals aged 12 years and older. Refer to the Ministry of Health COVID-19 Vaccine Guidance document (Version 1.0 or as current) for further information and possible symptoms.¹⁰
- Individuals with a proven severe allergic reaction (e.g. anaphylaxis) to injectable therapy not related to a component of the COVID-19 vaccines (e.g. other intramuscular, intravenous or subcutaneous vaccines or therapies) may be routinely vaccinated with COVID-19 vaccines with a 30 minute post-vaccination observation period.²
- Individuals with a suspected but unproven allergy to a vaccine component (e.g. PEG) may be routinely vaccinated with COVID-19 vaccines with a 30 minute post-vaccination observation period.²

- Individuals with a history of allergy not related to a component of the COVID-19 vaccines or other injectable therapy (e.g., foods, oral drugs, insect venom or environmental allergens) may be routinely vaccinated with Covid-19 vaccines with a 15 minute post-vaccination observation period.²
- In individuals with bleeding disorders, the condition should be managed prior to immunization to minimize the risk of bleeding. Individuals receiving long-term anticoagulation are not considered to be at higher risk of bleeding complications following immunization and may be safely immunized without discontinuation of their anticoagulation therapy.²
- Symptoms, either current or displayed recently, of chest pain or shortness of breath:
 - Vaccine should not be offered to persons displaying current or recent history of chest pain or shortness of breath.
 - Persons displaying current or recent history of chest pain or shortness of breath should consult with a health care provider prior to vaccination and/or if symptoms are severe, should be directed to the emergency department or instructed to call 911.¹⁰
- Vaccination of individuals who may be currently infected with SARS-CoV-2 is not known to have a detrimental effect on the illness. However, vaccination should be deferred in symptomatic individuals with confirmed or suspected SARS-CoV-2 infection, or those with respiratory symptoms, to minimize the risk of COVID-19 transmission at an immunization clinic/venue. If any person is identified with symptoms on arrival at the venue, they should not be immunized and should be instructed to seek medical and public health advice and follow current local public health measures.²
- As a precautionary measure and considering the need to be able to monitor for COVID-19 vaccine adverse events without potential confounding from symptoms of COVID-19 or other co-existing illnesses, people should wait until all symptoms of an acute illness are resolved before vaccinating with a COVID-19 vaccine.²

Additional Special Considerations and Booster Dose Eligibility

- After reviewing the published literature and evidence sought from manufacturers of COVID-19 vaccines, NACI approved changes to the guidance to indicate that vaccination with COVID-19 vaccines may take place at any time before, after or at the same visit as the TST or IGRA test.⁶
- For the most current information related to specific populations (e.g. pregnancy and breastfeeding, Individuals previously infected with SARS-CoV-2, Persons with an autoimmune condition, Immunocompromised persons, Travellers, Persons new to Canada), please refer to the Government of Canada COVID-19 vaccine: Canadian Immunization Guide (last date modified: 2022-09-09 or as current).⁵

Obtaining Consent

Appropriate informed consent must be obtained as per the *Guidelines for Implementing the Order/Delegated Procedure* and the client must NOT have any contraindications.

Guidelines for Implementing the Order/Delegated Procedure

Nurses and nursing students are expected to prepare and administer vaccine(s) to clients in a safe, effective, and ethical manner, and to be accountable for their actions. Nurses and nursing students are expected to meet the College of Nurses of Ontario's [Medication Practice Standard](#) by performing all the administration and verification steps to minimize the chance of error.

Nurses and nursing students are expected to consult with their Manager or designate if unable to perform their nursing duties in a safe and effective manner.

Responsibilities of the nurse and nursing student who implement the intervention include the following:

- Clarify that informed consent has been obtained
- Assess the client to determine whether the specific client conditions and any situational circumstances identified in the directive have been met
- Know the risks to the client of implementing the Order/Delegated Procedure
- Possess the knowledge, skills and judgement required to implement the directive safely
- Know the predictability of the outcomes of the intervention
- Determine whether management of the possible outcomes is within their practice; if so whether they are competent to provide management of outcomes and if not, whether the appropriate resources are available to assist as required.

Administration of vaccine must be in accordance with the manufacturers' product information monograph and the Government of Canada COVID-19 vaccine: Canadian Immunization Guide.^{1,2}

At all clinics where immunizing agents are administered, the _____ (AGENCY DESIGNATE) will be available for consult.

Informed Consent

To obtain informed consent, the nurse and/or nursing student must ensure:

- The purpose of the immunization, desired outcome, risks, and benefits are reviewed.
- The client has read and understood the information on the appropriate COVID-19 Vaccine Information Sheet(s). If the client cannot read the sheet, it should be read or explained to them. The nurse must ensure the client has no contraindications to the vaccine as specified in the "Contraindications" section and query regarding current health (e.g., immunosuppression) and precautions.
- The client has had the opportunity to ask questions about the vaccine and had these questions answered to their satisfaction; and

- The client has provided informed written or verbal consent to the nurse administering vaccine.

Capacity to provide consent

A client can give consent if they understand the information that is being presented to them and appreciate the foreseeable consequences of their decision or lack of decision.

When in doubt about informed consent or a medical condition of the client, do not administer the vaccine and consult with the _____ (AGENCY DESIGNATE).

Spikevax™ Bivalent COVID-19 Vaccine (Moderna)

COVID-19 Booster Dose Recommendations^{1,10}

Vaccination	Dose	Presentation	Dose Volume	Intervals
Booster Dose	50 mcg	Bivalent Moderna 0.10mg/mL (Royal Blue Vial Cap Green Label)	0.5mL	For population-specific booster dose recommendations including dosing, intervals and recommended vaccine, refer to the Ministry of Health document COVID-19 Vaccine Guidance document (Version 1.0 September 9, 2022 or as current). ¹⁰

Preparation and Administration of the Spikevax COVID-19 Vaccine

For injection technique, follow the _____ (NAMED AGENCY) procedure in conjunction with the following resources below.

For basic information on vaccine administration, refer to the Ministry of Health [COVID-19 Vaccine Guidance](#) document (Version 1.0 September 9, 2022 or as current).¹⁰

For step-by-step directions for administration, refer to the [Spikevax™ Bivalent COVID-19 Vaccine \(Moderna\) product monograph](#) (September 1, 2022 or as current).¹

Storage Requirements and Additional Doses from Vaccine Vials

For detailed information on storage requirements, refer to the [Ministry of Health Chapter 2: Storage and Handling of Moderna COVID-19 Vaccines](#) (Version 2: September 9, 2022, or as current)¹¹ document.

Documentation and Communication

Documentation of administered doses into the COVax database post-vaccination should be followed in conjunction with the _____ (NAMED AGENCY) documentation procedure, and the College of Nurses of Ontario's Documentation Practice Standard.⁴

Adverse Reactions

Non-Serious Adverse Reactions

The following side effects have been reported as being common or very common after Spikevax™ COVID-19 Vaccine (Moderna):¹

- Pain at the injection site
- Tiredness
- Headache
- Muscle ache and stiffness
- Chills
- Fever
- Swelling or redness at the injection site
- Nausea and/or vomiting
- Enlarged lymph nodes
- Hypoesthesia (decreased sense of touch or sensation, numbness) or paraesthesia (tingling, itching or pricking sensation)
- Dizziness

Post-Vaccination Observation

Recommendations for the post-vaccination observation period for other vaccines during the pandemic, such as for influenza vaccine, should continue to be followed. Therefore, advise clients to remain in the post-vaccination waiting area for the length of time as recommended in the Ministry of Health [COVID-19 Vaccine Guidance](#) document (Version 1.0 September 9, 2022 or as current).¹⁰

Serious Adverse Reactions

In situations of suspected hypersensitivity or non-anaphylactic allergy to COVID-19 vaccine components, the (AGENCY DESIGNATE) shall be consulted prior to vaccination, to determine the appropriateness of the vaccination, the option of vaccination in a controlled setting and/or the use of an extended post-vaccination observation period.

As per section 38 of the *Health Protection and Promotion Act*³, those administering vaccines should ensure that the vaccine recipients are advised to immediately call their doctor/nurse practitioner or go to the nearest hospital emergency department if they develop a reaction that could be related to the vaccine.³ Specifically, the client should be monitored for:

- Hives
- Swelling of the mouth and throat
- Altered level of consciousness/serious drowsiness
- Trouble breathing, hoarseness or wheezing
- High fever (over 40°C or 104°F)
- Convulsions (seizures)
- Other serious reactions (e.g., “pins and needles” or numbness)¹⁰

If a reaction as described above occurs while at (NAMED AGENCY) clinic, refer to the *Medical Directive for Anaphylaxis Management*.

Adverse events/reactions must be documented and reported to the Renfrew County and District Health Unit.

Approving Physician(s)/Authorizer(s):

<p>NAME:</p> <p>SIGNATURE:</p>	<p>DATE: YYYY/MM/DD</p>
---------------------------------------	-------------------------

Appendix A: Components of the Spikevax™ Bivalent COVID-19 Vaccine (Moderna) and its Packaging

For a comprehensive list of components in the vaccine and packaging, please consult the product leaflet or information contained within the Spikevax™ Bivalent COVID-19 Vaccine (Moderna) product monograph.¹

Potential non-medicinal ingredients in the vaccine or its container known to cause type 1 hypersensitivity reactions ranging from mild cutaneous reactions to anaphylaxis includes the following (which may not be a complete list):

- polyethylene glycol (PEG)₂

The full list of non-medical ingredients for multidose vial (2.5 mL) and multidose vial (5 mL) is as follows:¹

- Acetic acid
- Cholesterol
- DSPC (1,2-distearoyl-sn-glycero-3- phosphocholine)
- Lipid SM-102
- PEG2000-DMG (1,2-dimyristoyl-racglycerol,methoxy-polyethyleneglycol)
- Sodium acetate trihydrate
- Sucrose
- Trometamol
- Trometamol hydrochloride
- Water for injection

Spikevax™ Bivalent COVID-19 Vaccine (Moderna) does not contain any preservatives, antibiotics, adjuvants, or human- or animal-derived materials. The glass vials have a stopper which does not contain natural rubber latex.¹

REFERENCES

1. Product Monograph- Spikevax Bivalent COVID-19 Vaccine (Date Revised: September 1, 2022 or as current): [spikevax-bivalent-en.pdf \(canada.ca\)](#)
2. Government of Canada COVID-19 vaccine: Canadian Immunization Guide (Last Date Modified: September 9, 2022 or as current): <https://www.canada.ca/en/public-health/services/publications/healthy-living/canadian-immunization-guide-part-4-active-vaccines/page-26-covid-19-vaccine.html>
3. Health Protection and Promotion Act: <https://www.ontario.ca/laws/statute/90h07#BK45>
4. College of Nurses of Ontario. Documentation, revised 2008. Toronto, ON: 2019. Available from: http://www.cno.org/globalassets/docs/prac/41001_documentation.pdf
5. Government of Canada COVID-19 vaccine: Canadian Immunization Guide- Vaccination of specific populations (Last Date Modified: September 9, 2022 or as current): <https://www.canada.ca/en/public-health/services/publications/healthy-living/canadian-immunization-guide-part-4-active-vaccines/page-26-covid-19-vaccine.html#a6>
6. Government of Canada COVID-19 vaccine: Canadian Immunization Guide- Other considerations (Last Date Modified: September 9, 2022 or as current): <https://www.canada.ca/en/public-health/services/publications/healthy-living/canadian-immunization-guide-part-4-active-vaccines/page-26-covid-19-vaccine.html#a11>
7. Ministry of Health Document: [Chapter 2: Storage and Handling of Moderna COVID-19 Vaccines](#) (Version 2: September 9, 2022, or as current).
8. Government of Canada, COVID-19 vaccine guide for youth and adults (12 years and over): Managing COVID-19 vaccine administration errors or deviations (last date modified: 2022-02-03 or as current): <https://www.canada.ca/en/public-health/services/diseases/2019-novel-coronavirus-infection/guidance-documents/quick-reference-guide-covid-19-vaccines/managing-administration-errors-deviations.html>
9. College of Nurses of Ontario's *Medication Practice Standard* (Last Updated January 2019 or as current): https://www.cno.org/globalassets/docs/prac/41007_medication.pdf
10. Ministry of Health Document: [COVID-19 Vaccine Guidance](#), Version 1.0 September 9, 2022 (or as current).
11. NACI's Recommendations on the use of bivalent Omicron-containing mRNA COVID-19 Vaccines. September 1 2022: [NACI Recommendations on the use of bivalent Omicron-containing mRNA COVID-19 vaccines \(canada.ca\)](#)