


Management of Tick Bites and Investigation of Early Localized Lyme Disease

Please contact us at evidence@ontariohealth.ca or **1-877-280-8538** if you have any questions or feedback about this clinical guidance document.



Box 1. Clinical Manifestations of Early Localized Lyme Disease: Erythema Migrans Rashes



Additional images of typical and atypical rashes are available on [Health Canada's website](#); under "Clinical manifestations," please see "Erythema migrans rash."

Note: People with darker skin tones may present with a bruise-like rash.

Box 2. Prevalence of Symptoms in Patients Presenting With Possible Early Localized Lyme Disease[#]

- Erythema migrans rash (typical or atypical) ~70%
- Fatigue 54%
- Myalgia 44%
- Headache 42%
- Fever/chills 39%
- Stiff neck 35%
- Decreased appetite 26%

[#]As a disease of public health significance, Lyme disease is reportable in Ontario under the Health Protection and Promotion Act, R.S.O. 1990, c. H.7.

Box 4. Areas of Risk for Lyme Disease

- The risk of acquiring Lyme disease varies across geographical regions. Please click to see the risks in [Ontario, Canada](#), and the [United States](#)
- In Europe, the areas of highest risk are in Central and Eastern Europe, but infected ticks have also been found in Southern Scandinavia and up to the northern Mediterranean region


Box 5. Post-Exposure Prophylaxis

The risk of developing Lyme disease following a tick bite by an infected tick is between 1% and 3%. In Ontario, the prevalence of infected ticks varies by geographic region. In many instances, it is reasonable to adopt the "wait and see" approach and treat patients if they develop symptoms compatible with Lyme disease. Counsel patients to watch for the development of early signs and symptoms for 30 days, and advise patients that other tick-borne infections may result in signs or symptoms too.


Based on the best available evidence, post-exposure prophylaxis can be considered if these four criteria are met:

- The tick was attached > 24 hours
- The tick was removed within the past 72 hours
- The tick was acquired in an area with a prevalence of ticks infected with *Borrelia burgdorferi* > 20% (e.g., Rouge National Urban Park and Morningside Park in the Greater Toronto Area,

Box 3. Blacklegged Ticks at Various Stages and Safe Tick Removal



For more images, please go to: [Centers for Disease Control and Prevention](#).



For instructions, please see [Centers for Disease Control and Prevention](#).

Brighton, Kingston and surrounding areas, Thousand Islands, Brockville, Perth-Smiths Falls and surrounding areas, Ottawa and surrounding areas, Rondeau Provincial Park in Morpeth, and Pinery Provincial Park in Grand Bend*)

4. Doxycycline is not contraindicated. (Doxycycline is contraindicated for pregnant or lactating people and those with an allergy. There is insufficient evidence for the prophylactic use of other medications, such as amoxicillin)

Recommended treatment for post-exposure prophylaxis:

Adults: 1 dose of doxycycline 200 mg, by mouth

Children < 18 years of age: 1 dose of doxycycline 200 mg dose or 4 mg/kg (up to a maximum dose of 200mg), by mouth

*Note: This is not a comprehensive list of higher-risk areas in Ontario. Infectivity rate is not uniformly collected and updated, and therefore post-exposure prophylaxis decisions are sometimes made based on risk-benefit discussions with patients.

For more information, please refer to the [Ontario Lyme Disease Map](#).

References

- American Academy of Pediatrics Committee on Infectious Diseases. Lyme disease (Lyme Borreliosis, *Borrelia burgdorferi* sensu lato infection). In: Kimberlin DW, Brady MT, Jackson MA, editors. Red Book (2018): report of the Committee on Infectious Diseases: American Academy of Pediatrics; 2018.
- Cameron DJ, Johnson LB, Maloney EL. Evidence assessments and guideline recommendations in Lyme disease: the clinical management of known tick bites, erythema migrans rashes and persistent disease. *Expert Rev Anti Infect Ther*. 2014;12(9):1103–35.
- Canadian Paediatric Society (CPS). Lyme disease in Canada: focus on children [Internet]. Ottawa (ON): The Society; 2020 [cited 2022 Dec]. Available from: <https://cps.ca/documents/position/lyme-disease-children>
- Centre for Effective Practice (CEP). Early Lyme disease management in primary care [Internet]. Toronto (ON): The Centre; 2020 [cited 2022 Dec]. Available from: https://cep.health/media/uploaded/CEP_EarlyLymeDisease_Provider_2020.pdf
- Committee on Infectious Diseases American Academy of Pediatrics, Kimberlin DW, Barnett ED, Lynfield R, Sawyer MH. Lyme disease (Lyme Borreliosis, *Borrelia burgdorferi* sensu lato infection). Red Book: 2021–2024 report of the Committee on Infectious Diseases: American Academy of Pediatrics; 2021.
- Hatchette T, Lindsay R, on behalf of the Lyme Disease Diagnostic Working Group. Modified two-tiered testing algorithm for Lyme disease serology: the Canadian context. *Can Commun Dis Rep*. 2020;46(5):125–31.
- Pegalajar-Jurado A, Schriefer ME, Welch RJ, Couturier MR, MacKenzie T, Clark RJ, et al. Evaluation

- of Modified Two-Tiered Testing Algorithms for Lyme Disease Laboratory Diagnosis Using Well-Characterized Serum Samples. *J Clin Microbiol*. 2018;56(8):e01943–17.
- Lantos PM, Rumbaugh J, Bockenstedt LK, Falck-Ytter YT, Aguero-Rosenfeld ME, Auwaerter PG, et al. Clinical practice guidelines by the Infectious Diseases Society of America (IDSA), American Academy of Neurology (AAN), and American College of Rheumatology (ACR): 2020 guidelines for the prevention, diagnosis and treatment of Lyme disease. *Clin Infect Dis*. 2020;72(1):e1–e48.
- Meissner HC, Steere AC. Management of pediatric Lyme disease: updates from 2020 Lyme guidelines. *Pediatrics*. 2022;149(3). Aguero-Rosenfeld ME, Wang G, Schwartz I, Wormser GP. Diagnosis of Lyme Borreliosis. *Clin Microbiol Rev*. 2005;18(3):484–509.
- Nadelman RB. Erythema migrans. *Infect Dis Clin North Am*. 2015;29(2):211–39.
- National Institute for Health and Care Excellence (NICE). Lyme disease [Internet]. United Kingdom: The Institute; 2018 [cited 2022 Dec]. Available from: <https://www.nice.org.uk/guidance/ng95>
- National Institute for Health and Care Excellence (NICE). Lyme disease: diagnosis and management. [D] Evidence review for the management of erythema migrans [Internet]. United Kingdom: The Institute; 2018 [cited 2022 Dec]. Available from: <https://www.nice.org.uk/guidance/ng95/evidence/d-management-of-erythema-migrans-pdf-4792271010>
- Wormser GP, Dattwyler RJ, Shapiro ED, Halperin JJ, Steere AC, Klempner MS, et al. The clinical assessment, treatment, and prevention of Lyme disease, human granulocytic anaplasmosis, and babesiosis: clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis*. 2006;43(9):1089–134.

Box 6. Laboratory Testing

- Laboratory testing is generally not indicated for asymptomatic patients
- Serological testing may not yield positive results during early localized Lyme disease, so management should not be based on serological testing results during this phase
- Antibiotic treatment in early disease may reduce seroconversion; testing should not be used to monitor treatment outcome
- Following exposure to *Borrelia burgdorferi*, immunoglobulin M (IgM) antibodies are detected within 2–4 weeks, and IgG antibodies within 4–6 weeks
- As of April 1, 2023, Public Health Ontario uses a modified two-tiered testing (MTTT) algorithm to maximize sensitivity and specificity (see Box 7)
- For serological testing, please complete the [requisition](#) fully and submit it, along with samples, to a public health laboratory for testing
- If European Lyme disease is suspected based on the patient's travel history, please order serology testing

Box 7. Sensitivity of Serological (Modified Two-Tier) Testing[†] in Patients With Lyme Disease^{6,7}

Erythema migrans, acute phase (early localized disease)	58%
Erythema migrans, convalescence phase [‡] (early localized disease)	76%
Neurological involvement (early disseminated disease)	100%
Arthritis (late disseminated disease)	100%

[†]The MTTT algorithm is based on serum sample initially tested using IgG/IgM enzyme-linked immunosorbent assay (ELISA) using a whole cell lysate (tier 1). If results of tier 1 ELISA results are reactive/indeterminate, sample is further tested using second (tier 2) IgG/IgM ELISA assay targeting specifically VlsE1 and pepC10 antigens.

[‡]Following antibiotic treatment.

Box 8. Recommendations for Treatment of Patients With Early Localized Lyme Disease

Drugs	Dosage for Adults	Dosage for Children
Preferred		
Doxycycline*	100 mg twice a day for 10–21 days [†] Contraindicated for pregnant or lactating people	For children < 18 years of age: 4 mg/kg, orally divided into 2 doses (maximum 200 mg/day) for 10–21 days [†]
Amoxicillin	500 mg three times a day for 14–21 days	For children < 18 years of age: 50 mg/kg/day orally, divided into 3 equal doses per day, maximum of 500 mg per dose for 14–21 days
Cefuroxime	500 mg twice per day for 14–21 days	For children > 8 years of age: 30 mg/kg/day divided in 2 doses (maximum 500 mg/dose) for 14–21 days
For Allergy or Intolerance [‡]		
Azithromycin	500 mg/day for 7–17 days	For children < 18 years of age: 10 mg/kg/day, orally, once daily for 7–17 days
Clarithromycin	500 mg twice a day for 14–21 days Relatively contraindicated in pregnant people	For children > 8 years of age: 7.5 mg/kg twice a day (maximum 500 mg/day) for 14–21 days

[†]Doxycycline is considered to be the preferred antibiotic treatment option by some guidelines for early Lyme disease (erythema migrans) in both children and adults based on its ability to treat potential extracutaneous manifestations of infection (particularly neurological involvement) and potential coinfection or infection with *Anaplasma phagocytophilum* (anaplasmosis) or *Ehrlichia muris*-like agent.^{5,9,12}

[‡]Recent guidelines,^{3,5,8} including one by the Canadian Paediatric Society, recommend a 10-day treatment duration with doxycycline in children of all ages and adults. A 2018 guideline¹⁰ recommends a 21-day treatment duration with doxycycline in children over 9 years of age and adults based on concerns with low cure rates and a lack of clear evidence for shorter courses. In addition, a longer course may be reassuring for people being treated for early Lyme disease who continue to have symptoms and the evidence suggests that adverse event rates were not increased for longer courses.¹²

[‡]Patients treated with macrolides should be closely monitored to ensure resolution of clinical symptoms as macrolides are less effective.